

# Measurements of Turbulence in the Upper Layer with AUTOSUB

Thomas Osborn  
Olin Hall  
The Johns Hopkins University  
3400 N Charles Street  
Baltimore, MD 21218-2681  
phone (410) 516-7039 fax (410) 516-7933 email [osborn@jhu.edu](mailto:osborn@jhu.edu)

Award Number: N000149910088

## LONG-TERM GOALS

The long term goal is to understand the dynamics of the upper layer of the ocean, specifically the roles of turbulence, Langmuir circulation and large scale vortices.

## OBJECTIVES

- 1) To measure the turbulence, velocity and temperature fields in the upper layer in order to delineate the processes responsible for the heat and momentum fluxes
- 2) To develop the technical capability to operate from a wide variety of AUV and manned submersible vehicles

## APPROACH

Heat, momentum and gas transfer between the ocean and the atmosphere involve wave breaking, turbulence and bubble generation, mixing by Langmuir circulation and shear induced eddies; all these processes are poorly understood (Thorpe et. al., 2002). These processes control the dynamics of much of the forcing of the upper layer of the ocean.

Measurements from the USS Dolphin showed enhanced turbulent dissipation above 'Law of the Wall' scaling with acoustically detected bubble clouds in the upper 4m of the water column during 5-9 m/s winds off San Diego (Osborn et. al., 1992). More detailed measurements by Agrawal et. al. (1992) show the enhanced dissipation extends to a depth  $z \approx 10^5 u_*^2 / g$  where  $u_*$  is the friction velocity for the water and  $g$  the acceleration of gravity. Terray et. al. (1996) suggest that the significant wave height,  $H_s$  and the wave age,  $c / u_*$ , are relevant scaling parameters ( $c$  is the phase speed of the peak of the wave frequency spectrum). Another parameter they use is  $c'$ , where the rate of energy input from the wind,  $F$ , is  $F = c' \tau$ . However, work by Annis and Moum (1995) and Gemmrich and Farmer (1999a & b) suggest that the parameters  $H_s$ ,  $c'$ ,  $u_*$ , and  $z$  are not sufficient to scale the turbulence.

The surface turbulence affects the near surface bubble distribution and the transfer of gases and heat. Below this layer, Langmuir circulation (with vortices roughly aligned with the wind direction) contributes to the transport and distribution. Another large scale process is the shear induced eddies identified by Thorpe and Hall (1980) in Loch Ness. These vortices arise from a shear instability of the mean flow in the upper layer and thus their vorticity is aligned with the vorticity of the mean shear. In

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2003</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2003 to 00-00-2003</b>	
4. TITLE AND SUBTITLE <b>Measurements of Turbulence in the Upper Layer with AUTOSUB</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>The Johns Hopkins University,,3400 N Charles Street,,Baltimore,,MD, 21218</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

Loch Ness, the mean shear was in the direction of the wind and the vortices aligned perpendicularly to the wind. There they generated temperature ramps that produced more significant thermal features than those from the Langmuir circulation. The temperature signal from the ramps was observed by moving parallel to the wind direction and the temperature signal from the Langmuir circulation was observed by moving perpendicularly to the wind.

My approach is to perform turbulence measurements with airfoil probes for velocity fluctuations in conjunction with high resolution temperature and temperature gradient data from fast thermistors. These data are combined with data from other investigators in order to determine the interaction of the turbulence with breaking waves, bubbles, Langmuir circulation, and large scale eddies. We have used AUTOSUB in the past and are arranging access to that vehicle again in 2005. For the past 2 years we have used the submersible Forel in Lake Geneva.

## **WORK COMPLETED**

Professors Steve Thorpe, David Farmer and I collected a unique data set with AUTOSUB in April 2000 in the upper layer of the ocean. The system included: Thorpe's forward and sideways looking sonar (Thorpe and Hall, 1983) to document bubble clouds, Langmuir circulation, and breaking waves; Farmer's resonator system (Farmer et. al., 1998) to determine bubble size distribution; and my turbulence package to measure the turbulent dissipation and wave statistics. This year has seen the publication of the second paper (Thorpe et. al., 2003) on the relation between the turbulence, Langmuir circulation, bubbles and breaking waves from measurements in a Scottish loch. This paper tests (to the ability of the data) eight of the basic assumptions supporting models for bubble distributions and air-sea gas exchange. The assumption seems generally valid below  $1.5H_s$  in winds around 11.5 m/s. The paper concludes by drawing "attention to several aspects of upper ocean dynamics for which data are not yet sufficient to construct realistic and quantitative models of transfer processes involving Langmuir circulation."

In November 2001 and again in October 2002, Dr Ulrich Lemmin (Laboratoire de Recherches Hydrauliques of the Ecole Polytechnique Fédérale de Lausanne), Steve Thorpe and I used the submarine FOREL in Lake Geneva to attempt to collect data in the mixed layer during a strong wind regime. Past measurements in Lake Geneva by Dr. Lemmin and Prof. Thorpe have focused on breaking progressive internal waves during stratification and the plume formation and boundary mixing during convective winter cooling.

## **RESULTS**

The turbulence package we used on the FOREL is the same system used in the AUTOSUB measurements. It has nine channels of data: two shear probes (one measuring the cross stream horizontal velocity fluctuations and the other the vertical velocity component), temperature, temperature derivative, three accelerometers (for pitch, roll and heave), pressure and pressure derivative. The shear data are used to calculate the rate of dissipation of kinetic energy in 1 m horizontal bins. The temperature and temperature derivative data are combined and digitally filtered (Osborn et. al., 1992, using the method of Mudge and Lueck, 1994) to produce high resolution temperature traces with resolution of millidegrees. The pressure data from the turbulence package can be processed to produce a high resolution time series to estimate wave spectra (Osborn et. al., 1992). This allows us to estimate the significant wave height and period of the dominant waves. One data

channel is used to record signal pulses that are recorded concurrently on the thermistor array data in order to lock the two times series together unambiguously in time.

The data from both field test have been through preliminary processing to calculate time series of dissipation, temperature and temperature gradient variance. Preliminary integration of the thermistors array with the turbulence data has been accomplished and final integration of the data sets is planned for November 2003.



***Figure 1. The submarine Forel with sensors in the deployed position before a dive on Lake Geneva in October 2002. The turbulence package is mounted in foam rubber in the brown tube along the port side of the submarine. The thermistor array is shown in the deployed position with thermistors and Nortek 3-D acoustic current meters. For launch, transit and recovery, the turbulence package is retracted about 2 meters and the thermistor array is folded onto the upper deck. For October 2002 the thermistors array was modified (as shown) to include horizontal as well as the previous (November 2001) vertical spacing of the sensors. This array appears to be the largest practical for use on the submarine.***

## **IMPACT/APPLICATIONS**

The development of the capability to sample turbulence in the surface and near surface regimes of the upper ocean in difficult weather conditions is vital to understanding the important and complex processes that occur there.

## TRANSITIONS

The technology for turbulence measurements and interpretation that has been developed by the PI through ONR funding during the last 25 years is now well established in the oceanography community.

## REFERENCES

- Agrawal, Y. C., E. A. Terray, M. A. Donelan, P. A. Hwang, A. J. Williams III, W. M. Drennan, K. K. Kahma, and S. A. Kitagorodski 1992 Enhanced dissipation of kinetic energy beneath surface waves. *Nature*, **359**, 219-220.
- Anis, A., and J.N. Moum 1995 Surface wave-turbulence interactions. Scaling  $\varepsilon(z)$  near the sea surface. *J. Phys. Oceanogr.*, **25**, 2025-2045.
- Farmer, D. M., S. Vagle, and A. O. Booth 1998 A free-flooding acoustical resonator for measurement of bubble size distributions. *Atmos. Ocean. Tech.*, **15**, 1132-1146.
- Gemmrich, J.R., and D.M. Farmer 1999a Near surface turbulence and thermal structure in a wind driven sea. *J. Phys. Oceanogr.*, **29**, 480-499.
- Gemmrich, J.R., and D.M. Farmer 1999b Observations of the scale and occurrence of breaking surface waves. *J. Phys. Oceanogr.*, **29**, 2595-2606.
- Mudge, T.D. and R.G. Lueck, 1994. Digital signal processing to enhance oceanographic observations, *J. Atmos. Oceanogr. Techn.*, **11**, 825-836.
- Osborn, T., D.M. Farmer, S. Vagle, S. Thorpe, M. Cure, 1992. Measurements of bubble plumes and turbulence from a submarine. *Atmosphere - Ocean* **30**, 419-440.
- Terray, E. A., M. A. Donelan, Y. C. Agrawal, W. M. Drennan, K. K. Kahma, A. J. Williams III, P. A. Hwang, and S. A. Kitagorodski 1996 Estimates of kinetic energy dissipation under breaking waves. *J. Phys. Oceanogr.*, **26**, 792-807.
- Thorpe, S.A., 1995. Dynamical processes of transfer at the sea surface. *Prog. Oceanogr.* **35**, 315-352.
- Thorpe, S.A., 1996. Interactions between internal waves and boundary layer vortices. *Journal of Physical Oceanography*, **27**, 62-71.
- Thorpe, S. A. and A. J. Hall 1980 The mixing layer of Loch Ness, *J. Fluid Mech.*, **101**, (4), 687-703.
- Thorpe, S. A. and A. J. Hall 1983 The characteristics of breaking waves, bubble clouds and near-surface currents observed using side-scan sonar. *Continental Shelf Research*, **1**, (4), 353-384.
- Thorpe, S.A., T. R. Osborn, J. F. E. Jackson, A. J. Hall, and R. G. Lueck 2002 Measurements of turbulence in the upper ocean mixing layer using Autosub. *J. Phys. Oceanogr.*, **33**, 122-145.
- Thorpe, S.A., T. R. Osborn, D.M. Farmer, and S. Vagle 2003 Bubble clouds and Langmuir circulation: observations and models. *J. Phys. Oceanogr.*, **33**, 122-145.

## PUBLICATIONS

Thorpe, S.A., T. R. Osborn, J. F. E. Jackson, A. J. Hall, and R. G. Lueck 2002 Measurements of turbulence in the upper ocean mixing layer using Autosub. *J. Phys. Oceanogr.*, **33**, 122-145.

Thorpe, S.A., T. R. Osborn, D.M. Farmer, and S. Vagle 2003 Bubble clouds and Langmuir circulation: observations and models. *J. Phys. Oceanogr.*, **33**, 122-145.